

Lawrence Berkeley National Laboratory

Recent Work

Title

VARIATIONAL FORMULATION OF COVARIANT EIKONAL THEORY FOR VECTOR WAVES

Permalink

<https://escholarship.org/uc/item/61f3h90q>

Authors

Kaufman, A.N.

Ye, H.

Hui, Y.

Publication Date

1986-10-01

2



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Accelerator & Fusion Research Division

RECEIVED
LIBRARY
BERKELEY LABORATORY

NOV 18 1986

LIBRARY AND
DOCUMENTS SECTION

Presented at the Fourth Symposium on Energy
Engineering Sciences, Argonne, IL, May 7-9, 1986;
and submitted to Physics Letters A


VARIATIONAL FORMULATION OF COVARIANT EIKONAL THEORY FOR VECTOR WAVES

A.N. Kaufman, H. Ye, and Y. Hui

October 1986

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.*



LBL-21472 Rev.
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

VARIATIONAL FORMULATION OF COVARIANT EIKONAL THEORY FOR VECTOR WAVES*

Allan N. Kaufman, Huanchun Ye, and Yukkei Hui

Lawrence Berkeley Laboratory and Physics Department
University of California
Berkeley, CA 94720

ABSTRACT

The eikonal theory of wave propagation is developed by means of a Lorentz-covariant variational principle, involving functions defined on the natural eight-dimensional phase space of rays. The wave field is a four-vector representing the electromagnetic potential, while the medium is represented by an anisotropic, dispersive nonuniform dielectric tensor $D^{\mu\nu}(k,x)$. The eikonal expansion yields, to lowest order, the Hamiltonian ray equations, which define the Lagrangian manifold $k(x)$, and the wave-action conservation law, which determines the wave-amplitude transport along the rays. The first-order contribution to the variational principle yields a concise expression for the transport of the polarization phase. The symmetry between k -space and x -space allows for a simple implementation of the Maslov transform, which avoids the difficulties of caustic singularities.

*Supported by U.S. DOE Contract # DE-AC03-76SF00098.

Wave propagation in various media is often well represented by the eikonal theory, also known as ray optics, geometrical optics, and WKB [1]. In this paper, we present a variational approach, leading to some new results, as well as concise expressions for old results. While our immediate motivation is electromagnetic wave propagation in magnetized plasma [2], we expect that our methods are applicable to other media, such as elastic waves [3].

Our emphasis here is on the vector character of the wave field [4], which propagates through an anisotropic medium. We also emphasize the phase-space concept, as a way of avoiding the singularities of caustics.

In addition, it is convenient to utilize a covariant formulation [5], treating space and time on an equal footing; this is the key to conciseness. We represent the electromagnetic wave field by the four-potential $A_\mu(x)$, with $A_\mu = (\underline{A}, -\phi)$, and $x = (\underline{x}, t)$. (We set $c = 1$).

The wave equation for linear dissipationless propagation can be expressed as an integral equation of the form

$$\int d^4x'' D^{\mu\nu}(x', x'') A_\nu(x'') = 0. \quad (1)$$

Here $D^{\mu\nu}(x', x'')$ represents the (in general) anisotropic dispersion tensor of the medium, as a two-point kernel. We consider it as given; methods for its derivation are discussed in Ref. [6]. The variational equation below requires that it be symmetric: $D^{\nu\mu}(x', x'') = D^{\mu\nu}(x'', x')$; asymmetry represents dissipation, which must be treated by other methods. Nonlinearity can also be included, by allowing the medium to respond "ponderomotively" to the wave [6].

We introduce the quadratic action functional:

$$S(A) = \frac{1}{2} \int d^4 x' \int d^4 x'' A_\mu(x') D^{\mu\nu}(x', x'') A_\nu(x''). \quad (2)$$

The requirement that S be stationary with respect to arbitrary variations of the four-potential field is equivalent to Eq. (1). It is advantageous to introduce the phase-space concept as early as possible. We define the dispersion tensor as a function on phase space (k, x) :

$$D^{\mu\nu}(k, x) = \int d^4 s D^{\mu\nu}(x' = x + s/2, x'' = x - s/2) \exp(-ik \cdot s) \quad (3)$$

It is thus the Fourier transform of $D^{\mu\nu}(x', x'')$ with respect to the two-point separation $s = x' - x''$. The wave four vector $k = (\underline{k}, -\omega)$, together with $x = (\underline{x}, t)$, coordinatizes the eight-dimensional phase space (k, x) of the rays.

Analogously, we introduce the bilinear Wigner tensor:

$$A_{\mu\nu}^2(k, x) = \int d^4 s A_\mu(x + s/2) A_\nu(x - s/2) \exp(-ik \cdot s), \quad (4)$$

which likewise lives on the phase space. In terms of these, the action functional reads

$$S(A) = \frac{1}{2} \iint D^{\mu\nu}(k, x) A_{\nu\mu}(k, x), \quad (5)$$

where

$$\iint = \iint d^4 x d^4 k / (2\pi)^4.$$

We see that k -space and x -space enter (5) on an equal footing. At each point (k, x) in this space, we represent the tensor $D^{\mu\nu}$ in terms of its local (orthonormal) eigenvectors e_α and (real) eigenvalues D_α :

$$D^{\mu\nu}(k, x) = \sum_{\alpha} D_{\alpha}(k, x) e_{\alpha}^{\mu}(k, x) e_{\alpha}^{\nu*}(k, x), \quad (6)$$

where

$$D^{\mu\nu}(k, x) e_{\nu}^{\alpha}(k, x) = D_{\alpha}(k, x) e_{\alpha}^{\mu}(k, x)$$

and

$$e_{\alpha}^{\mu*}(k, x) e_{\mu}^{\beta}(k, x) = \delta_{\alpha}^{\beta}.$$

(We assume that the eigenvalues are non-degenerate [7].) Substituting (6) into (5), we obtain

$$S(A) = \frac{1}{2} \sum_{\alpha} \iint D_{\alpha}(k, x) A_{\alpha}^2(k, x), \quad (7)$$

where

$$A_{\alpha}^2(k, x) = e_{\alpha}^{\nu*}(k, x) A_{\nu\mu}^2(k, x) e_{\alpha}^{\mu}(k, x) \quad (8)$$

is the projection of the Wigner tensor on the local polarization $e_{\alpha}(k, x)$, and $D_{\alpha}(k, x)$ is the scalar dispersion function.

We now assume that the wave field $A_{\mu}(x)$ can be expressed in eikonal form:

$$A_{\mu}(x) = a_{\mu}(x) \exp i\theta(x) + c.c., \quad (9)$$

where the phase $\theta(x)$ is real and has slowly varying first-derivative

$k_{\mu}(x) = \partial\theta(x)/\partial x^{\mu}$, while the amplitude $a_{\mu}(x)$ is complex and is slowly varying.

When we substitute (9) into (4), we obtain, to lowest order in the eikonal parameter,

$$A_{\mu\nu}^2(k, x) = (2\pi)^4 [\delta^4(k - \partial\theta/\partial x) a_{\mu}^*(x) a_{\nu}(x) + \delta^4(k + \partial\theta/\partial x) a_{\mu}(x) a_{\nu}^*(x)], \quad (10)$$

where we have discarded rapidly oscillating terms $\sim \exp 2i\theta(x)$, which vanish upon x -integration in (5).

Substituting (10) into (8), and performing the k -integration of (7), we obtain the action functional $S(A) = \int d^4x L(x)$ in terms of a Lagrangian density:

$$L(x) = \sum_{\alpha} D_{\alpha}(k = \partial\theta/\partial x, x) |a_{\alpha}|^2(x), \quad (11)$$

where

$$a_{\alpha}(x) = e_{\alpha}^{\mu}(k = \partial\theta/\partial x, x) a_{\mu}(x) \quad (12)$$

is the scalar projection of the amplitude on the polarization direction e_{α} .

The variation of the action S is now taken with respect to the amplitude $a_{\mu}(x)$ and the phase $\theta(x)$. The former variation yields

$$D_{\alpha}(k = \partial\theta/\partial x, x) a_{\alpha}(x) = 0, \quad (13)$$

for each α . For a non-trivial solution, we require $D_{\alpha} = 0$ for one polarization, denoted I , and allow $a_{\alpha} = 0$ for the other polarizations $\alpha \neq I$.

Thus we obtain the eikonal equation

$$D_I(k=\partial\theta/\partial x, x) = 0, \quad (14)$$

with the amplitude expressed as

$$a_\mu(x) = e_\mu^I(x) a_I(x). \quad (15)$$

The polarization

$$e_\mu^I(x) = e_\mu^I(k=\partial\theta/\partial x, x) \quad (16)$$

can absorb the complex phase of the amplitude vector field $a_\mu(x)$, allowing the scalar amplitude $a_I(x)$ to be real and positive.

The eikonal equation (14) is solved by Hamilton's method. In eight-dimensional phase space, the ray equations are

$$\frac{dx^\mu}{d\sigma} = - \frac{\partial D_I}{\partial k_\mu}, \quad \frac{dk_\mu}{d\sigma} = + \frac{\partial D_I}{\partial x^\mu}. \quad (17)$$

These Hamiltonian equations yield the family of ray orbits $[k(\sigma), x(\sigma)]$. With appropriate initial conditions, the rays generate a four-dimensional "Lagrangian submanifold" $k(x)$ [8]. The phase $\theta(x)$ is then obtained by integrating $k(\sigma)$ along a ray $x(\sigma)$.

On varying S with respect to the phase $\theta(x)$, we obtain the conservation law

$$\partial J^\mu(x) / \partial x^\mu = 0, \quad (18)$$

for the wave-action four-flux $J^\mu(x)$ [9], defined as

$$J^\mu(x) = - \partial L / \partial k_\mu = - [\partial D_I(k, x) / \partial k_\mu](x) a_I^2(x) = (dx^\mu / d\sigma) a_I^2(x). \quad (19)$$

The temporal component of $J^\mu(x)$ is the familiar wave-action density:

$$J(\underline{x}; t) = (\partial D_I / \partial \omega) a_I^2,$$

while the spatial part of $J^\mu(x)$ is the wave-action flux density $(\partial \omega / \partial \underline{k}) J$.

Thus (18) reads

$$\partial J(\underline{x}; t) / \partial t = - \nabla \cdot (J \partial \omega / \partial \underline{k});$$

this conservation law reflects the invariance of the action functional under a uniform phase shift $\theta(x) \rightarrow \theta(x) + C$.

On substituting expression (19) for $J^\mu(x)$ into the conservation law (18), we obtain the amplitude transport equation:

$$\frac{d}{d\sigma} \ln a_I^2 = \frac{\partial^2 D_I(k, x)}{\partial k_\mu \partial x^\mu} + \frac{\partial^2 D_I}{\partial k_\mu \partial k_\nu} \frac{\partial^2 \theta(x)}{\partial x^\mu \partial x^\nu} \quad (20)$$

The first term on the right represents the medium nonuniformity, while the second represents the divergence (or convergence) of a ray bundle.

The latter quantity is determined, in turn, by its transport relation:

$$\begin{aligned} \frac{d}{d\sigma} \frac{\partial^2 \theta}{\partial x^\mu \partial x^\nu} &= \frac{\partial^2 D_I}{\partial x^\mu \partial x^\nu} + \frac{\partial^2 D_I}{\partial x^\mu \partial k_\rho} \frac{\partial^2 \theta}{\partial x^\rho \partial x^\nu} + \frac{\partial^2 D_I}{\partial x^\nu \partial k_\rho} \frac{\partial^2 \theta}{\partial x^\rho \partial x^\mu} \\ &+ \frac{\partial^2 D_I}{\partial k_\rho \partial k_\lambda} \frac{\partial^2 \theta}{\partial x^\lambda \partial x^\mu} \frac{\partial^2 \theta}{\partial x^\rho \partial x^\nu} \end{aligned} \quad (21)$$

obtained by differentiating (14) twice with respect to x . This nonlinear equation for the ray divergence leads in general to a singularity in a finite distance, i.e., to a caustic. To avoid that singularity, we may use the Maslov transform [8].

Inserting (15) into (9), we have $A_\mu(x) = e_\mu^I(x) a_I(x) \exp i\theta(x) + \text{c.c.}$, with $\theta(x)$ determined by (17), $a_I(x)$ determined by (20), and $e_\mu(x)$ determined by (16) up to a complex phase factor. That is, the replacement $e_\alpha^H(k,x) \rightarrow e_\alpha^H(k,x) \exp i\psi(x)$ leaves (6) invariant, so that the polarization phase is not as yet determined.

For the polarization phase, it is necessary to expand the action functional to first order in the eikonal parameter. After some algebra, and using the zero-order equations, we find the first order Lagrangian to be

$$L'(x) = i a_I^2(x) (e_I \cdot de_I / d\sigma + \frac{1}{2} D^{\mu\nu} \{e_\nu^I, e_\mu^{I*}\}), \quad (22)$$

where $\{, \}$ represents the canonical Poisson bracket:

$$\{f, g\} = \frac{\partial f}{\partial x^\mu} \frac{\partial g}{\partial k_\mu} - \frac{\partial f}{\partial k_\mu} \frac{\partial g}{\partial x^\mu}.$$

On taking the variation of the action functional with respect to the amplitude, we now obtain the additional equation $L' = 0$, which yields the desired polarization phase transport:

$$e_I^* \cdot \frac{de_I}{d\sigma} = - \frac{1}{2} D^{\mu\nu} \{e_\nu^I, e_\mu^{I*}\}. \quad (23)$$

Note that this transport relation lives in phase space; i.e., it is not necessary to project onto x -space. On the other hand, the ray divergence equation (21) definitely refers to x -space.

We now proceed to the elimination of caustic singularities. Let us again substitute the Wigner tensor (10) into (8), obtaining the scalar field:

$$A_I^2(k, x) = (2\pi)^4 a_I^2(x) [\delta^4(k - \partial\theta/\partial x) + \delta^4(k + \partial\theta/\partial x)] \quad (24)$$

Integrating over an element of phase-space volume [see (7)], we have

$$d^4x d^4k A_I^2(k, x) / (2\pi)^4 = a_I^2(x) d^4x. \quad (25)$$

We see that $A_I^2(k, x)$ is the wave density in phase space, while $a_I^2(x)$ is the density in x-space. From (24), we see that A_I^2 is supported on the Lagrangian manifold $k = k(x) = \partial\theta(x)/\partial x$, i.e., $k_1 = \partial\theta(x)/\partial x^1$, and so on. At a caustic, $k(x)$ becomes double-valued, so that the x-representation of the eikonal breaks down.

The Maslov procedure [8] is a simple way to avoid the caustic singularity. One Fourier transforms the field $A_\mu(x)$ from (x^1, x^2, x^3, x^4) space, to (k_1, x^2, x^3, x^4) space, where the (x^1, k_1) pair is chosen to represent the singular direction in $\partial^2\theta/\partial x^\mu \partial x^\nu$. One can now make the eikonal assumption (9) in the new space, and the whole procedure outlined above carries through in the same way. Space does not permit a fuller discussion here. The Maslov transform is a special case of a more general approach, recently developed by Littlejohn [10-12]. His method involves a continuous transformation of coordinates in phase space, and is based on the wave-packet approximation.

The variational method discussed here allows for a treatment of wave angular momentum [13] transport, and its consequences. This work is in progress. Applications of the present formalism are under way.

ACKNOWLEDGMENTS

This research was supported by the Office of Basic Energy Sciences of the U.S. Department of Energy under Contract Number DE-AC0376SF00098. We have benefited from discussions with B. Boghosian, R. Littlejohn, S. McDonald, S. Omohundro, and P. Similon.

REFERENCES

1. I. BERNSTEIN and L. FRIEDLAND, "Geometric Optics in Space- and Time-Varying Plasma," in *Handbook of Plasma Physics*, Vol. I, p. 367 (R. Sagdeev and M. Rosenbluth, eds.; North Holland, Amsterdam, 1983).
2. S. McDONALD and A. KAUFMAN, "Weyl Representation for Electromagnetic Waves: The Wave Kinetic Equations," *Phys. Rev.* 32A, 1708 (1985).
3. J. ACHENBACH, A. GAUTESEN, and H. McMAKEN, *Ray Methods for Waves in Elastic Solids* (Pitman, Boston, 1982).
4. H. BERK and D. PFIRSCH, "WKB Method for Systems of Integral Equations," *J. Math. Phys.* 21, 2054 (1980).
5. A. KAUFMAN and D. HOLM, "The Lie-Transformed Vlasov Action Principle: Relativistically Covariant Wave Propagation and Self-Consistent Ponderomotive Effects," *Phys. Letters* 105A, 277 (1984).
6. S. McDONALD, C. GREBOGI, and A. KAUFMAN, "Locally Coupled Evolution of Wave and Particle Distribution in General Magnetoplasma Geometry," *Phys. Lett.* 111A, 19 (1985).
7. For the near degenerate case, see L. FRIEDLAND, "Renormalized Geometric Optics Description of Mode Conversion in Weakly Inhomogeneous Plasmas," *Phys. Fluids* 28, 3260 (1985).
8. V. MASLOV and M. FEDORIUK, "Semi-Classical Approximation in Quantum Mechanics" (Reidel, Dordrecht, 1981).
9. R. DEWAR, "Lagrangian Derivation of the Action-Conservation Theorem," *Astrophys. J.* 174, 301 (1972).
10. R. LITTLEJOHN, "Semiclassical Evolution of Wave Packets," *Physics Reports* 138, 193 (1986).
11. R. LITTLEJOHN, "Symplectically Invariant WKB Wave Functions," *Phys. Rev. Lett.* 54, 1742 (1985).
12. R. LITTLEJOHN, "Wave Packet Evolution and Quantization," *Phys. Rev. Lett.* 56, 2000 (1986).
13. D. SOPER, "Classical Field Theory" (Wiley, New York, 1976), Chapter 9.

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

*LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720*