

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

The Sudbury Neutrino Observatory

### Permalink

<https://escholarship.org/uc/item/7nz6t5k6>

### Authors

Norman, E B

Chan, Y D

Garcia, A

et al.

### Publication Date

1992-11-01



# Lawrence Berkeley Laboratory


UNIVERSITY OF CALIFORNIA

Presented at the APS Division of Particles and Fields Meeting,  
Batavia, IL, November 11-14, 1992, and to be published in the Proceedings

## The Sudbury Neutrino Observatory

E.B. Norman, Y.D. Chan, A. Garcia, K.T. Lesko, A.R. Smith, R.G. Stokstad, I. Zliven,  
H.C. Evans, G.T. Ewan, A. Hallin, H.W. Lee, J.R. Leslie, J.D. MacArthur, H.B. Mak,  
A.B. McDonald, W. McLatchie, B.C. Robertson, P. Skensved, B. Sur, E. Bonvin,  
E.D. Earle, D. Hepburn, G.M. Milton, I. Blevis, W.F. Davidson, C.K. Hargrove, H. Mes,  
M. Shatkay, D. Sinclair, A.L. Carter, B. Hollebone, D. Kessler, P. Jagam, J. Law,  
R.W. Ollerhead, J.J. Simpson, J.X. Wang, J. Bigu, E.D. Hallman, R.U. Haq, C. Virtue,  
S. Gil, C. Waltham, M.M. Lowry, E.W. Beier, W. Frati, M. Newcomer, R. Van Berg,  
T.J. Bowles, P. Doe, M.M. Fowler, A. Hime, F. McGirt, R.G.H. Robertson,  
T.C. Spencer, D.J. Vieira, J.B. Wilhemy, J.F. Wilkerson, J.M. Wouters, N.W. Tanner,  
N.A. Jelley, J.C. Barton, G. Doucas, E.W. Hooper, A.B. Knox, M.E. Moorhead,  
M. Omori, P.T. Trent, and D.L. Wark

November 1992



Prepared for the U.S. Department of Energy under Contract Number DE-AC03-76SF00098

REFERENCE COPY	1	LBL-33304
Does Not Circulate	1	Copy 1
Bldg. 50 Library.	1	

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

# The Sudbury Neutrino Observatory

E. B. Norman, Y.D. Chan, A. Garcia, K.T. Lesko, A.R. Smith, R.G. Stokstad, I. Zliten  
Nuclear Science Division, Lawrence Berkeley Laboratory  
University of California, Berkeley, California 94720, USA

H.C. Evans, G.T. Ewan, A. Hallin, H.W. Lee, J.R. Leslie, J.D. MacArthur,  
H.B. Mak, A.B. McDonald, W. McLatchie, B.C. Robertson, P. Skensved, B. Sur  
Physics Department, Queen's University  
Kingston, Ontario K7L 9N6, Canada

E. Bonvin, E.D. Earle, D. Hepburn, G.M. Milton  
AECL Research, Chalk River Laboratories  
Chalk River, Ontario K0J 1J0, Canada

I. Blevis, W.F. Davidson, C.K. Hargrove, H. Mes, M. Shatkay, D. Sinclair  
Centre for Research in Particle Physics, Carleton University  
Ottawa, Ontario K1S 5B6, Canada

A.L. Carter, B. Hollebhone, D. Kessler  
Physics Department, Carleton University  
Ottawa, Ontario K1S 5B6, Canada

P. Jagam, J. Law, R.W. Ollerhead, J.J. Simpson, J.X. Wang  
Physics Department, University of Guelph  
Guelph, Ontario N1G 2W1, Canada

J. Bigu, E.D. Hallman, R. U. Haq, C. Virtue  
Physics Department, Laurentian University  
Sudbury, Ontario P3E 2C6, Canada

S. Gil, C. Waltham  
Physics Department, University of British Columbia  
Vancouver, British Columbia V6T 1W5, Canada

M.M. Lowry  
Physics Department, Princeton University  
Princeton, NJ 08544 U.S.A.

E.W. Beier, W. Frati, M. Newcomer, R. Van Berg  
Physics Department University of Pennsylvania  
Philadelphia, PA 19104 U.S.A.

T. J. Bowles, P. Doe, M.M. Fowler, A. Hime, F. McGirt, R.G.H. Robertson,  
T.C. Spencer, D.J. Vieira, J.B. Wilhemy, J.F. Wilkerson, J.M. Wouters  
Physics Division, Los Alamos National Laboratory  
Los Alamos, NM 87545 U.S.A.

N.W. Tanner, N.A. Jelley, J.C. Barton, G. Doucas, E.W. Hooper,  
A.B. Knox, M.E. Moorhead, M. Omori, P.T. Trent, D.L. Wark  
Physics Department, University of Oxford  
Oxford, OX1 3RH, U.K.

This work was supported by the Director, Office of Energy Research Division  
of Nuclear Physics of the Office of High Energy and Nuclear Physics of the  
U.S. Department of Energy under Contract DE-AC03-76SF00098

## 1. Introduction

Two experiments now in progress have reported measurements of the flux of high energy neutrinos from the Sun. Since about 1970, Davis and his co-workers<sup>1</sup> have been using a  $^{37}\text{Cl}$ -based detector to measure the  $^7\text{Be}$  and  $^8\text{B}$  solar neutrino flux and have found it to be at least a factor of three lower than that predicted by the Standard Solar Model<sup>2</sup> (SSM). The Kamiokande collaboration<sup>3</sup> has been taking data since 1986 using a large light-water Cerenkov detector and have confirmed that the flux is about two times lower than predicted. Recent results from the SAGE<sup>4</sup> and GALLEX<sup>5</sup> gallium-based detectors show that there is also a deficit of the low energy pp solar neutrinos. These discrepancies between experiment and theory could arise because of inadequacies in the theoretical models of solar energy generation or because of previously unobserved properties of neutrinos. The Sudbury Neutrino Observatory (SNO) will provide the information necessary to decide which of these solutions to the "solar neutrino problem" is correct.

## 2. The SNO Detector

The Sudbury Neutrino Observatory will consist of a 1000 tonne heavy water ( $\text{D}_2\text{O}$ ) Cerenkov detector that is designed to measure the flux, energy spectrum, and direction of neutrinos from the Sun and from such other sources as supernovae. It is presently under construction in a very low background environment 2000 meters underground in the Creighton mine near Sudbury, Ontario, Canada. This is an operating nickel mine owned by INCO, Ltd. The  $\text{D}_2\text{O}$  used in the detector will be on loan from Atomic Energy of Canada Limited.

The basic measurements that will be made with the SNO detector are:

- 1) the flux and energy spectrum of electron-type neutrinos reaching the Earth, and
- 2) the total flux of all neutrino types above an energy of 2.2 MeV.

With these two measurements, it will be possible to :

- 1) determine if neutrino oscillations occur, and
- 2) independently test solar models by determining the production rate of high energy electron-type neutrinos in the solar core.

The SNO detector utilizes three complementary neutrino interactions with the heavy water.

- 1) The neutrino-electron elastic scattering (ES) reaction:  $\nu_x + e^- \rightarrow \nu_x + e^-$ .

The observed signal in the detector is the Cerenkov light produced by the recoiling electron. This is the primary detection mechanism for light water detectors such as the Kamioka detector. It is sensitive to all neutrino types, but is dominated by the electron neutrino. The recoiling electrons from the ES reaction are strongly forward peaked and give excellent directional information. However, information about the energy spectrum of the neutrinos is more difficult to extract because of averaging over the outgoing neutrinos.

- 2) The charged current (CC) reaction:  $\nu_e + d \rightarrow p + p + e^-$  ( $Q = -1.44 \text{ MeV}$ ).

This reaction of the electron-type neutrino on the deuteron is unique to the SNO detector. It has a relatively large cross section for  $^8\text{B}$  neutrinos and would produce about 10 events per day for one third of the SSM flux. This is greater than fifty times more sensitive than existing solar neutrino experiments. The electron energy is  $E_e = E_\nu - 1.44$

MeV and the energy resolution is approximately 20%. This reaction gives good spectral information on the  $^8\text{B}$  neutrinos and thus provides good sensitivity to the MSW effect. This reaction will also identify electron neutrinos from the initial burst of a supernova.

3) The neutral current (NC) reaction:  $\nu_x + d \rightarrow \nu_x' + p + n$  ( $Q = -2.2$  MeV).

This reaction can be observed by the detection of the gamma rays resulting from the subsequent neutron capture or by a neutron detector array in the heavy water. This reaction is sensitive to all types of neutrinos equally and would be used to measure the total flux of neutrinos above the threshold energy of 2.2 MeV. The expected counting rate for the full SSM is approximately 10 per day. This will give a direct measure of the total solar  $^8\text{B}$  neutrino production independent of neutrino oscillation effects. It will also detect all types of neutrinos from supernovae explosions.

The  $\text{D}_2\text{O}$  target of the SNO detector will be contained in a transparent spherical acrylic vessel with a diameter of 12 m and a wall thickness of 5 cm. Approximately 2.5 m outside the acrylic vessel, there will be about 9600 photomultipliers (PMTs) with 20-cm diameters uniformly arranged and held in place by a geodesic support structure. A reflector is mounted in front of each PMT to increase the light collection to yield a total effective photocathode coverage of approximately 60%. The PMT array is sensitive to Cerenkov radiation produced by relativistic electrons and other particles in the central regions of the detector. The acrylic vessel, PMTs, and the support structure are immersed in 7300 tonnes of ultrapure  $\text{H}_2\text{O}$ . This reduces the background in the heavy water from radioactive impurities in the rock walls and in the detector components. The cavity which will house the detector is barrel shaped with a diameter of 22 m and a height of 30 m. The excavation of the cavity is expected to be completed early in 1993. The PMT support structure and acrylic vessel will then be assembled and installed underground. The initial water fill is scheduled to begin near the end of 1994, with a view toward completing commissioning tests and starting to take data in 1995. The data taking sequence that will be used in the experiment is still under discussion. It may begin with a  $\text{H}_2\text{O}$  fill of the acrylic vessel followed by a  $\text{D}_2\text{O}$  fill. The NC measurement may then be made either by adding 2.5 tonnes of  $\text{NaCl}$  to the  $\text{D}_2\text{O}$  (in order to raise the energy of the capture gamma rays) or through the use of discrete neutron detectors (such as  $^3\text{He}$  counters) installed in the  $\text{D}_2\text{O}$ .

### 3. Acknowledgements

This research is supported by the National Science and Engineering Research Council, National Research Council, Ministry of Industry, Science and Technology, and Northern Ontario Heritage Fund Corporation, all in Canada, the Department of Energy in the United States, and the Science and Engineering Research Council in the United Kingdom. The loan of heavy water from Atomic Energy of Canada Limited is greatly appreciated.

### 4. References

1. R. Davis, Jr. *et al.*, *Phys. Rev. Lett.* **20** (1968) 1205.  
J.K. Rowley *et al.*, in *Solar Neutrinos and Astronomy*, eds. M. L. Cherry, W. A. Fowler, and K. Lande (AIP, New York, 1985).
2. J. N. Bahcall, *Neutrino Astrophysics* (Cambridge Univ. Press, Cambridge, 1989).
3. K. Hirata *et al.*, *Phys. Rev. Lett.* **65** (1990) 1297.
4. A. Abuzov *et al.*, *Phys. Rev. Lett.* **67** (1991) 3332.
5. P. Anselmann *et al.*, *Phys. Lett.* **B285** (1992) 376.

LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
TECHNICAL INFORMATION DEPARTMENT  
BERKELEY, CALIFORNIA 94720